

CLINICAL TRIALS AND OBSERVATIONS

Venetoclax-rituximab with or without bendamustine vs bendamustine-rituximab in relapsed/refractory follicular lymphoma

Pier Luigi Zinzani,¹ Ian W. Flinn,² Sam L. S. Yuen,³ Max S. Topp,⁴ Chiara Rusconi,⁵ Isabelle Fleury,⁶ Katell Le Dû,⁷ Christopher Arthur,⁸ Barbara Pro,⁹ Giuseppe Gritti,¹⁰ Michael Crump,¹¹ Adam Petrich,¹² Divya Samineni,¹³ Arijit Sinha,¹⁴ Elizabeth A. Punnoose,¹³ Edith Szafer-Glusman,¹³ Nathalie Spielewoy,¹⁵ Mehrdad Mobasher,¹³ Kathryn Humphrey,¹⁴ Martin Kornacker,¹⁵ and Wolfgang Hiddemann¹⁶

¹Institute of Hematology "L. e A. Seràgnoli," University of Bologna, Bologna, Italy; ²Sarah Cannon Research Institute/Tennessee Oncology, Nashville, TN; ³Department of Haematology, Calvary Mater Newcastle, Waratah, NSW, Australia; ⁴Medizinische Klinik und Poliklinik II, Universitätsklinikum Würzburg, Würzburg, Germany; ⁵Division of Hematology, Niguarda Hospital, Milan, Italy; ⁶Department of Hematology, Maisonneuve-Rosemont Hospital and University of Montreal, Montreal, QC, Canada; ⁷Hematology Department, Clinique Victor Hugo, Le Mans, France; ⁸Department of Haematology, Royal North Shore Hospital, St Leonards, NSW, Australia; ⁹Robert H. Lurie Comprehensive Cancer Center, Chicago, IL; ¹⁰Hematology and BMT Unit, Ospedale Papa Giovanni XXIII, Bergamo, Italy; ¹¹Princess Margaret Cancer Centre, University of Toronto, Toronto, ON, Canada; ¹²AbbVie, North Chicago, IL; ¹³Genentech, Inc., South San Francisco, CA; ¹⁴Roche Products Limited, Welwyn Garden City, United Kingdom; ¹⁵F. Hoffmann-La Roche Ltd, Basel, Switzerland; and ¹⁶Department of Medicine III, University of Munich, Munich, Germany

KEY POINTS

- Chemotherapy-free VEN + R had modest efficacy and acceptable toxicity in relapsed/refractory disease warranting further study.
- VEN + BR led to increased toxicity; optimized chemotherapeutic dosing and/or combinations remain to be explored.

This open-label phase 2 study (CONTRALTO) assessed the safety and efficacy of BCL-2 inhibitor venetoclax (VEN) plus rituximab (R), and VEN plus bendamustine (B) and R, vs B + R (BR) alone in relapsed/refractory (R/R) follicular lymphoma. Patients in the chemotherapy-free arm (arm A: VEN + R) received VEN 800 mg/d plus R 375 mg/m² on days 1, 8, 15, and 22 of cycle 1 and day 1 of cycles 4, 6, 8, 10, and 12. After a safety run-in with VEN 600 mg, patients in the chemotherapy-containing cohort were randomized to either VEN + BR (arm B; VEN 800 mg/d for 1 year + 6 cycles of BR [B 90 mg/m² on days 1 and 2 and R 375 mg/m² on day 1]) or 6 cycles of BR (arm C). Overall, 163 patients were analyzed (9 in the safety run-in and 52, 51, and 51 in arms A, B, and C, respectively). Complete metabolic/complete response rates were 17% (arm A), 75% (arm B), and 69% (arm C). Of patients in arm B, only 61% received ≥90% of the planned B dose vs 96% of patients in arm C. More frequent hematologic toxicity resulted in more reduced dosing/treatment discontinuation in arm B vs arm C. Rates of grade 3/4 adverse events were 51.9%, 93.9%, and 60.0% in arms A, B, and C, respectively. VEN + BR led to increased toxicity and lower dose intensity of BR than in arm C, but efficacy was similar. Optimizing dose and schedule to maintain BR dose intensity

may improve efficacy and tolerability of VEN + BR, while VEN + R data warrant further study. This study was registered at www.clinicaltrials.gov as #NCT02187861. (*Blood*. 2020;136(23):2628-2637)


Introduction

Follicular lymphoma (FL) is typically treated by combining an anti-CD20 antibody with chemotherapy, which improves response rates, progression-free survival (PFS), and overall survival compared with chemotherapy alone.¹⁻⁵ However, many patients relapse repeatedly, with progressively increasing resistance to therapy.^{6,7} Use of targeted agents such as BCL-2 inhibitors may enhance antitumor therapy by acting as chemosensitizers.⁸⁻¹⁰

Venetoclax (VEN) is a highly selective, potent oral BCL-2 inhibitor, approved globally in multiple indications, including use in chronic lymphocytic leukemia (CLL) patients who have received ≥1

previous therapy, either in combination with rituximab (R) or as monotherapy in Europe.¹¹ In the United States, approval is for the treatment of adult patients with CLL or small lymphocytic lymphoma and patients with previously untreated acute myeloid leukemia who are ineligible for intensive chemotherapy, in combination with hypomethylating agents or cytarabine.¹² Pre-clinical data in CLL and non-Hodgkin lymphoma suggest that VEN + R or VEN + bendamustine and R (BR) may improve response compared with R or chemotherapy alone.^{10,13} Early clinical data also support the safety and efficacy of VEN in FL as monotherapy or combined with BR.^{13,14} The present study (CONTRALTO; NCT02187861) assessed VEN + R and VEN + BR vs BR alone in patients with relapsed or refractory FL.

Figure 1. Dosing schedule by arm and time on study. Arm A (chemotherapy-free): VEN (800 mg daily) + R (375 mg/m²). Arm B (chemotherapy containing): VEN (800 mg daily) + B (90 mg/m²) + R (375 mg/m²). Arm C (chemotherapy containing): B (90 mg/m²) + R (375 mg/m²). *R administered on days 1, 8, 15, and 22. Safety run-in arm (not displayed above) consisted of 9 patients that received VEN (600 mg daily for 1 year) + B (90 mg/m² on days 1 and 2 of each 28-day cycle) + R (375 mg/m² on day 1 of each cycle). D, day.



Methods

Study design and treatment

This open-label, international, multicenter phase 2 study comprised a safety run-in plus 3 treatment arms. Patients were enrolled into a chemotherapy-free (arm A: VEN + R) or chemotherapy-containing cohort at the investigator's (INV's) discretion. In the chemotherapy-containing cohort, patients were randomized 1:1 to arm B (VEN + BR) or arm C (BR only; Figure 1) using stratified permuted block randomization following a safety run-in (first 9 patients enrolled into the chemotherapy-containing cohort). Stratification was according to duration of response (DOR) to prior therapy (≤ 12 months/ > 12 months) and disease burden (high/low), according to modified Groupe d'Etude des Lymphomes Folliculaires criteria.¹⁵ Patients enrolled to the safety run-in received VEN 600 mg orally daily during 6 28-day cycles of standard BR (B 90 mg/m² IV on days 1 and 2 and R 375 mg/m² IV on day 1) and then continued VEN alone for 1 year. Following a safety review of the safety run-in and data from another phase 1 study¹³ by an internal monitoring committee and scientific oversight committee, the VEN dose for arm B during randomization was selected as 800 mg, given orally daily for 1 year, plus 6 28-day cycles of standard BR. Patients in arm C received 6 28-day cycles of standard BR. Patients chosen for arm A received VEN 800 mg orally daily + R (375 mg/m² IV on days 1, 8, 15, and 22 of cycle 1 and day 1 of cycles 4, 6, 8, 10, and 12) for 1 year (Figure 1). VEN treatment commenced on day 1 cycle 1 in study arms A and B.

All patients received prophylaxis for tumor lysis syndrome (TLS) before first dose of VEN + R or VEN + BR, including hydration and a uric acid-reducing agent.¹⁶

Institutional review boards/ethics committees at all participating institutions approved the protocol. The study was conducted in accordance with International Conference on Harmonization guidelines, including Good Clinical Practice, and the principles of the Declaration of Helsinki.¹⁷ All patients provided written informed consent.

Patient population

Eligible patients were aged ≥ 18 years with histologically confirmed FL (grade 1-3a); adequate coagulation, renal, and hepatic function; and had ≥ 1 prior FL therapy. Response duration of prior B, if received, had to be > 1 year in patients receiving chemotherapy. Refractoriness to last line of prior treatment was defined as progressive disease (PD) within 6 months of last dose or no response to treatment if no PD was reported and resistance

to R was defined as lack of response to or progression within 6 months of R-containing therapy. Exclusion criteria included primary central nervous system lymphoma, live vaccines within 28 days, and chemotherapy within 28 days of initiating study treatment.

Study objectives and end points

The primary efficacy objectives were to evaluate the activity of VEN + BR compared with BR and VEN + R, as measured by 18-fluorodeoxyglucose positron emission tomography (PET)-assessed complete metabolic response (CMR) rate 6 to 8 weeks after C6D1 (primary response assessment [PRA]), defined by the independent review committee (IRC) and in accordance with Lugano 2014 criteria.¹⁸

Secondary efficacy end points included PET-assessed CMR rate at PRA, as defined by the INV, PET-assessed CMR at 1 year from C1D1 (IRC and INV), computed tomography (CT)-assessed complete response (CR) at PRA and at 1 year from C1D1 (IRC and INV), overall response rate (ORR; IRC and INV), INV-assessed DOR (from first partial response or CR until PD or death), and PFS.

Safety objectives were to evaluate the safety and tolerability of (1) VEN 600 mg + BR in the safety run-in, (2) VEN+BR vs BR, and (3) VEN + R. Safety end points included the incidence, severity, and outcome of adverse events (AEs), changes in clinical laboratory results, and maintenance of relative dose intensity of B. Patients were followed for safety outcomes until at least 30 days after the last dose of VEN or B, at least 90 days since the last dose of R, and until study discontinuation or termination.

PET and CT imaging

Imaging at PRA and at 1 year included PET and CT scan with oral and IV contrast. Combined PET/CT scans were encouraged.

Biomarker analysis

BCL-2 immunohistochemistry (IHC) and BCL-2 fluorescence in situ hybridization (FISH) were performed, and minimal residual disease (MRD) negativity (per million count below the limit of detection) was evaluated in peripheral blood. See supplemental Appendix (available on the *Blood* Web site) for detailed methodology.

Data analysis and statistical methods

Efficacy analyses were conducted on the intent-to-treat population (enrolled patients). The safety population comprised patients who received ≥ 1 dose of any study treatment. The pharmacokinetic (PK)-evaluable population comprised patients who received ≥ 1 dose of VEN with ≥ 1 PK parameter estimated using noncompartmental analysis.

CMR rate and INV-based PET and CT response rates at PRA (and corresponding 95% confidence intervals [CIs]) were constructed according to the Clopper-Pearson method.¹⁹ The difference in CMR rates between arms B and C was calculated and the approximate 95% CI was determined using the Wald method.²⁰ Time-to-event end points were estimated using Kaplan-Meier methodology.²¹ Cox regression and logistic regression were used for exploratory analyses of potential biomarker effects on response probability and risk of progression.

Dose intensities were calculated as (actual dose/actual time)/(planned dose/planned time) \times 100%. For patients who discontinued treatment due to AEs, actual time was calculated to the time of planned treatment. For all other patients, including those with PD or those who died, actual time was calculated to the time of actual treatment.

Following an amendment to the statistical analysis plan due to the number of scans outside the prespecified window and missing data for the analysis period, an additional analysis of all available response assessments was conducted, and an extended analysis period of 42 to 65 weeks was used (see supplemental Appendix for details). Here, we focus on INV-assessed response assessment according to this updated statistical analysis plan for a more comprehensive efficacy assessment.

Expected sample size was ~ 156 patients (50 patients in each arm plus at least 6 patients in the safety run-in). Under a binomial distribution assumption, which was based on a 40% to 50% CMR rate as per historical BR data,^{22,23} this would allow 95% CIs for estimation of CMR to have a margin of error $\sim 15\%$.

Unless otherwise stated, baseline was defined as the last value obtained before the first dose of study drug.

Statistical analyses were performed using SAS version 9.4.

Results

Patient population

Patients (N = 163) were enrolled at 57 sites across 8 countries and followed between 13 November 2014 and 16 March 2018 (last patient, last visit), of whom 9 entered the safety run-in, 52 entered arm A (VEN + R), 51 entered arm B (VEN + BR), and 51 entered arm C (BR) (Figure 2). Three patients (2 in arm B and 1 in arm C) did not receive study treatment; therefore, the safety population comprised 160 patients.

Patient demographics and baseline clinical characteristics are summarized in Table 1. Demographics for patients from the safety run-in and patients who completed VEN + BR therapy are shown in supplemental Tables 1 and 2, respectively. At baseline, 120 out of 137 patients (87.6%) had BCL-2–positive disease (IHC

score $\geq 2/3$), with similar proportions noted across the study arms (Table 1). In 86 out of 106 patients with available data (81.1%), BCL-2 rearrangement was present, with similar proportions across all study arms.

Ten patients (4 in arm A, 5 in arm B, and 1 in arm C) had received an autologous stem cell transplant (supplemental Table 3). One patient in each arm was initially treated for diffuse large B-cell lymphoma and relapsed with diagnosis of FL.

A greater proportion of VEN + R patients (arm A; 92.3%) discontinued the study compared with VEN + BR (arm B; 62.7%) or BR (arm C: 62.7%), primarily due to a higher percentage with PD (80.8% vs 37.3% and 43.2%, respectively). Four patients discontinued the study due to an AE (1, 2, and 1 in arms A, B, and C, respectively), and 6 died (3, 1, and 2 in arms A, B, and C, respectively). Overall, 80 out of 103 patients with baseline samples (77.7%) tested MRD positive and represented the MRD-evaluable cohort.

Efficacy

Per the updated statistical analysis plan described above in "Methods," INV-assessed CMR/CR rates by PET + CT at the PRA were 75% (95% CI, 60.37 to 85.67) for arm B and 69% (95% CI, 54.11 to 80.89) for arm C (difference: 5.88 [95% CI, -11.59 to 23.35], $P = .51$), and at 1 year were 43% (95% CI, 29.35 to 57.75) and 51% (95% CI, 36.60 to 65.25), respectively (difference: -7.84 [95% CI, -27.16 to 11.47], $P = .43$) (Table 2). ORR by PET + CT was 84% for both arms B and C at the PRA and 49% and 57%, respectively, at 1 year. In comparison, the INV-assessed ORR by PET + CT was lower with VEN + R (arm A) at the PRA (35%) and at 1 year (27%) (Table 2); 17% of patients in arm A reached CMR at the PRA. Response rates were higher in a subgroup of 26 patients in arm A with nonrefractory FL vs those with refractory FL (ORR as best overall response [BOR] 54% vs 19%, respectively). In a subgroup of patients who were R refractory, INV-assessed CMR/CR rates by PET + CT at the PRA were 64% (95% CI, 35.14 to 87.24) for arm B, 67% (95% CI, 43.03 to 85.41) for arm C, and 11.8% (95% CI, 1.46 to 36.44), respectively. CMR as BOR by INV-assessed PET + CT was 25% in arm A (35% in nonrefractory and 15% in refractory FL patients) (Table 2). No significant differences in response between patients with refractory and nonrefractory FL were observed in arms B and C. IRC- and INV-assessed response rates in all study arms using PET + CT scans at the PRA and after 1 year of treatment were consistent (supplemental Table 4). INV-assessed responses at 1 year in the 19 patients who completed arm B (VEN + BR) are listed in supplemental Table 5.

With median follow-up of 18 months in arms B and C, the hazard ratio (HR) for DOR for arm B vs C was 0.69 (95% CI, 0.38 to 1.27; Figure 3A); INV-assessed median PFS was similar (HR, 0.69; 95% CI, 0.38 to 1.24) (Figure 3B). Median PFS was 6.6 months for arm A. Landmark analysis of PFS is shown in supplemental Figure 1.

Nearly all patients were positive by BCL-2 IHC and by BCL-2 FISH and negative by MCL-1 IHC (Table 1), limiting the ability to perform subgroup analysis for these biomarkers. PFS was not differentiated by BCL-XL IHC status (BCL-XL high: HR, 0.37 [95% CI, 0.11 to 1.2]; BCL-XL low: HR, 0.9 [95% CI, 0.43 to 1.9]). Analysis of response by expression of BCL-2 family genes showed no clear differences in response between arm A, and arm B vs C (supplemental Figure 2).




Figure 2. Patient flow. Rand, randomization.

In VEN + R responders (CMR/partial metabolic response as BOR), the rate of undetectable MRD at mid-induction (cycle 4) was 90% (9/10 MRD-evaluable patients), comparable with that of BR patients (100%; 7/7 MRD-evaluable patients).

Safety

Ninety-eight percent of patients in arm A and all patients in arms B and C had ≥ 1 AE (Table 3). Fifty-two patients had ≥ 1 serious AE, with the highest incidence in arm B (53.1%). A total of 103 patients (68.2%) had ≥ 1 grade 3/4 AE, most commonly neutropenia and thrombocytopenia.

In total, 41 patients (27.2%) received ≥ 1 course of granulocyte-colony stimulating factor for neutropenia prophylaxis: 3 (5.8%) patients in arm A, 21 (42.9%) in arm B, and 17 (34.0%) in arm C (supplemental Table 6). More patients in arm B received platelet transfusions (14.3%) compared with arm A (3.8%) and arm C (2.0%; supplemental Table 6).

Of 5 fatal AEs (Table 3), 3 were in arm A (pulmonary hemorrhage, colitis, myocardial infarction; all considered unrelated to study treatment); 1 in arm B (pneumonia, considered related to study treatment); and 1 in arm C (hypoxia in the context of pulmonary embolism, considered unrelated to study treatment).

No clinical TLS occurred. Four patients experienced grade 3/4 laboratory TLS: 1 (1.9%) in arm A and 3 (6.1%) in arm B. All patients were able to restart treatment.

AEs led to treatment discontinuation in 25 patients: 3 (5.8%) in arm A (2 VEN and 2 R discontinuations), 20 (40.8%) in arm B (17 VEN, 10 B, and 7 R discontinuations), and 2 (4.0%) in arm C (1 B and 1 B + R discontinuation). For details, see supplemental Table 7.

AE-related VEN dose modification or interruption occurred in 63 patients (20 in arm A and 43 in arm B), mostly due to neutropenia (n = 6) and diarrhea (n = 5) with VEN + R and neutropenia (n = 26) and thrombocytopenia (n = 18) with VEN + BR. Safety summaries for patients who completed VEN + BR and patients in the safety run-in are presented in supplemental Tables 8 and 9, respectively.

PK assessments

Ninety percent or more of the planned dose was received by 76.9% (VEN) and 90.4% (R) of patients in arm A; by 26.5% (VEN), 85.7% (R), and 61.2% (B) in arm B; and by 97.9% (R) and 95.8% (B) in arm C. Drug exposure data are shown in supplemental Table 10.

VEN PK parameters, available in 97 patients, were similar across arms A and B, suggesting no marked effect of BR or R co-administration (supplemental Table 11).

Discussion

CONTRALTO investigated the safety and efficacy of VEN in a chemotherapy-free regimen (VEN + R) and a B-containing regimen (VEN + BR) vs BR in patients with relapsed or refractory FL. VEN + R was associated with acceptable toxicity but modest activity. Most VEN + R-treated patients had advanced stage disease, were refractory to last treatment, or were heavily pretreated, and approximately one-third of them were refractory to R. However, responding patients had deep and sustained responses, similar to those receiving B-containing regimens. Patients receiving VEN + R who were nonrefractory to last treatment had superior response rates and MRD negativity compared with the total arm A population.

Table 1. Patient demographics and baseline clinical characteristics

Characteristic	Arm A: VEN + R (n = 52)	Arm B: VEN + BR (n = 51)	Arm C: BR (n = 51)
Median age, y (range)	63 (40-84)	66 (43-82)	61 (35-80)
Age \geq 65 y, n (%)	23 (44.2)	29 (56.9)	22 (43.1)
Male, n (%)	27 (51.9)	35 (68.6)	30 (58.8)
Lymph node \geq 10 cm, n (%)	5 (9.6)	4 (7.8)	7 (13.7)
Ann Arbor stage, n (%)	n = 50	n = 49	n = 51
I	2 (4.0)	5 (10.2)	4 (7.8)
II	4 (8.0)	8 (16.3)	10 (19.6)
III	9 (18.0)	13 (26.5)	7 (13.7)
IV	35 (70.0)	23 (46.9)	30 (58.8)
ECOG performance status, n (%)	n = 52	n = 48	n = 50
0	36 (69.2)	28 (58.3)	34 (68.0)
1	16 (30.8)	19 (39.6)	16 (32.0)
2	0	1 (2.1)	0
FL grade 3a, n (%)	n = 52 7 (13.5)	n = 50 7 (14.0)	n = 50 9 (18.0)
Bone marrow infiltration, n (%)	n = 51	n = 51	n = 49
Yes	18 (35.3)	19 (37.3)	13 (26.5)
No	32 (62.7)	31 (60.8)	35 (71.4)
Unknown	1 (2.0)	1 (2.0)	1 (2.0)
Extranodal involvement, n (%)			
Yes	31 (59.6)	30 (58.8)	27 (52.9)
No	21 (40.4)	21 (41.2)	24 (47.1)
Prior therapies, n			
Minimum to maximum	1-6	1-6	1-4
Median	3	3	2
Refractory* to last treatment, n (%)	26 (50.0)	19 (37.3)	23 (45.1)
Refractory to R, n (%)	17 (32.7)	14 (27.5)	21 (41.2)
Duration of prior therapy response, n (%)	n = 49	n = 51	n = 50
\leq 12 mo	29 (59.2)	22 (43.1)	26 (52.0)
>12 mo	20 (40.8)	29 (56.9)	24 (48.0)
Disease burden (GELF), n (%)	n = 52	n = 51	n = 51
Low	9 (17.3)	14 (27.5)	17 (33.3)
High	43 (82.7)	37 (72.5)	34 (66.7)
BCL-2 IHC evaluable, n (%)	n = 48	n = 46	n = 43
Negative (0-1)	6 (12.5)	4 (8.7)	7 (16.3)
Positive (2-3)	42 (87.5)	42 (91.3)	36 (83.7)
BCL-2 FISH evaluable, n (%)	n = 40	n = 33	n = 33
Negative	3 (7.5)	4 (12.1)	3 (9.1)
Positive	35 (87.5)	24 (72.7)	27 (81.8)
Undetermined	2 (5.0)	5 (15.2)	3 (9.1)
BCL-XL IHC evaluable, n (%)	n = 42	n = 46	n = 42
IHC score \geq 2,3	9 (21)	15 (33)	14 (33)
MCL-1 IHC evaluable, n (%)	n = 37	n = 38	n = 38
IHC score $\dagger \geq$ 2	0 (0)	0 (0)	2 (5)

ECOG, Eastern Cooperative Oncology Group; GELF, Groupe d'Etude des Lymphomes Folliculaires.

*Refractoriness to last line of prior treatment was defined as PD within 6 mo of last dose received or no response to treatment (if no PD was reported).

 \dagger No IHC scores \geq 3 were observed for MCL-1.

Table 2. Investigator-assessed response rates in the chemotherapy-free cohort (arm A) and chemotherapy-containing cohort (arms B and C) using PET + CT scan (intent-to-treat population) and BOR in arm A (refractory and nonrefractory patient subgroups)

N (%)	Arm A: VEN + R (n = 52)	Arm B: VEN + BR (n = 51)	Arm C: BR (n = 51)
Primary response assessment*			
ORR	18 (35)	43 (84)	43 (84)
CMR/CR	9 (17)	38 (75)	35 (69)
95% CI (Clopper-Pearson)	8.23, 30.33	60.37, 85.67	54.11, 80.89
Difference (95% CI)	NA	5.88 (−11.59 to 23.35), <i>P</i> = .51	
PMR/PR	9 (17)	5 (10)	8 (16)
NMR/SD	7 (14)	0	1 (2)
PMD/PD	23 (44)	2 (4)	5 (10)
Response data missing†	3 (6)	4 (8)	2 (4)
1 y follow-up‡			
ORR	14 (27)	25 (49)	29 (57)
CMR/CR	10 (19)	22 (43)	26 (51)
95% CI (Clopper-Pearson)	9.63, 32.53	29.35, 57.75	36.60, 65.25
Difference (95% CI)	NA	−7.84 (−27.16 to 11.47), <i>P</i> = .43	
PMR/PR	4 (8)	3 (6)	3 (6)
NMR/SD	3 (6)	0	0
PMD/PD	32 (62)	12 (24)	15 (29)
Response data missing‡	3 (6)	14 (28)	7 (14)
BOR, n (%)			
	Arm A: VEN + R (n = 52)	Arm A: VEN + R refractory§ (n = 26)	Arm A: VEN + R nonrefractory§ (n = 26)
ORR	19 (37)	5 (19)	14 (54)
CMR	13 (25)	4 (15)	9 (35)
PMR	6 (12)	1 (4)	5 (19)
NMR	6 (12)	4 (15)	2 (8)
PD	13 (25)	6 (23)	7 (27)
Response data missing†	14 (27)	11 (42)	3 (12)

NA, not applicable; NMR, no metabolic response; PMR, partial metabolic response; PR, partial response; SD, stable disease.

*Primary response evaluated from mid-induction visit to 10 weeks after C6D1 or 10 weeks after day 1 of the last cycle.

†Includes off study early for toxicity, off study early for not meeting inclusion/exclusion criteria, off study for death, withdrawal of consent, or unreadable scans. Responses were evaluated by PET/CT scan first; if missing, CT scan response was imputed. All patients with PD prior to or during the observation period were included as PD.

‡The 12-month responses were evaluated 42 to 65 weeks after C1D1.

§Refractoriness to last line of prior treatment was defined as PD within 6 months of last dose received or no responses to treatment (if no PD was reported).

Arms B and C showed similar CMR/CR rates and response durability; however, toxicity was higher with VEN + BR than with BR. More patients receiving VEN + BR than BR withdrew from study treatments or underwent dose modification, driven by hematologic and gastrointestinal toxicities. Only 27% of patients achieved $\geq 90\%$ VEN dose intensity in the VEN + BR arm, showing the limited tolerability of this regimen at the given dose and schedule. It is worth noting that patients in arm C on average were younger and had fewer prior lines of therapy compared with arm B. Indeed, the heavier pretreatment history of patients in arm B vs C may have had a compounding effect on the efficacy and tolerability/safety of VEN + BR. Despite the increased toxicity in arm B vs C, leading to lower BR exposure, similar efficacy outcomes were seen between the 2 arms. Further investigation is needed to determine the optimal dose and schedule of BR in combination with VEN for relapsed or refractory FL in order to maximize efficacy while minimizing toxicity. Indeed, the combination of VEN plus other chemotherapy regimens may offer a different safety profile and is worthy of further exploration.

The most important toxicities in the VEN-containing arms were grade 3/4 neutropenia and thrombocytopenia. The rate of neutropenia in the VEN + BR arm was similar to previously reported data for BR.²⁴ The rate of thrombocytopenia in the VEN + BR arm was notably higher than previously reported for BR.^{24,25} Incidences of gastrointestinal toxicities were consistent with the safety profile of VEN in other trials.^{14,26-29} TLS risk was mitigated by prophylaxis, and no cases of clinical TLS were reported. No new safety signals were reported with VEN treatment.

Limitations of the study design and conduct preclude precise conclusions on the efficacy of adding VEN to BR or R, but the benefit should not be dismissed. The dosing schedule used in this study was based on encouraging results from the phase 1 study,¹⁴ with doses as high as 1200 mg and also clearing the safety run-in. However, more toxicity was seen in the current study with the VEN + BR combination, and this led to increased dose modification. In the CAVALLI study, VEN was administered on a noncontinuous dosing schedule in combination with R plus cyclophosphamide, doxorubicin, vincristine, and prednisolone (CHOP) in diffuse large




Figure 3. Kaplan-Meier plot of DOR and PFS. DOR (A) and PFS (B) *Stratified by DOR prior therapy (≤ 12 months/ > 12 months) and disease burden (high/low), according to modified Groupe d'Etude des Lymphomes Folliculaires criteria.¹⁵

B-cell lymphoma and FL patients.^{30,31} On this schedule, VEN was associated with a higher rate of AEs; however, R-CHOP dose intensity was maintained at a similar rate to the historical comparator (R-CHOP arm of the GOYA study³²) with improved efficacy in the BCL-2–positive population.³¹ While comparisons are limited by different chemotherapy backbones, they highlight the potential of intermittent VEN administration over a longer treatment period to improve efficacy and tolerability of chemotherapy combination regimens. An important limitation of the protocol was the lack of mandatory guidance on dose reduction and discontinuation, and doses were modified at the INV's discretion at a higher rate than the historical comparator. Furthermore, VEN treatment was limited to 1 year, whereas targeted therapies have typically been administered until progression in other trials.

In conclusion, our data suggest that further investigation is warranted to confirm the clinical effects of adding VEN to

chemoimmunotherapy regimens or monoclonal antibodies in FL patients. In particular, optimization of VEN dosing and scheduling is needed, perhaps with noncontinuous dosing schedules being of particular interest in distinct populations. Several studies are now underway exploring VEN in various combination strategies in FL patients (NCT02956382, NCT03113422, NCT03135262, NCT02611323, and NCT02877550), with both continuous and noncontinuous schedules being assessed in first-line and relapsed/refractory settings. Notably, a phase 2 study is exploring obinutuzumab + B in combination with a noncontinuous dosing schedule of VEN (induction therapy) in previously untreated patients with high tumor burden FL (NCT03113422). The results of this study will help clarify how best to deliver VEN in B-containing regimens to maximize efficacy. The potential of VEN + R as a chemotherapy-free option for patients with nonrefractory FL also merits further exploration, particularly in earlier-line or frail populations who cannot tolerate chemotherapy. In addition, further

Table 3. Summary of safety (safety population)

	Arm A: VEN + R (n = 52)	Arm B: VEN + BR (n = 49)	Arm C: BR (n = 50)
Total number of AEs	425	936	502
Patients with ≥1 event, n (%)			
Any grade AE	51 (98.1)	49 (100.0)	50 (100.0)
Grade 3/4 AE	26 (50.0)	45 (91.9)	29 (58.0)
AE with fatal outcome (grade 5)	3 (5.8)	1 (2.0)	1 (2.0)
SAE	16 (30.8)	26 (53.1)	10 (20.0)
AE leading to dose modification/ interruption from any treatment	30 (57.7)	44 (89.8)	21 (42.0)
AE leading to VEN dose modification/ interruption	20 (38.5)	43 (87.8)	NA
AE leading to withdrawal from any treatment	3 (5.8)	20 (40.8)	2 (4.0)
AE leading to withdrawal of VEN	2 (3.8)	17 (34.7)	NA
All AEs occurring in ≥20% of patients in any treatment arm, n (%)			
Nausea	14 (26.9)	32 (65.3)	22 (44.0)
Neutropenia	14 (26.9)	30 (61.2)	17 (34.0)
Thrombocytopenia	7 (13.5)	28 (57.1)	8 (16.0)
Diarrhea	21 (40.4)	24 (49.0)	11 (22.0)
Vomiting	7 (13.5)	24 (49.0)	13 (26.0)
Fatigue	13 (25.0)	21 (42.9)	15 (30.0)
Anemia	3 (5.8)	19 (38.8)	7 (14.0)
Infusion-related reaction	18 (34.6)	10 (20.4)	7 (14.0)
Constipation	5 (9.6)	10 (20.4)	17 (34.0)
Cough	6 (11.5)	12 (24.5)	12 (24.0)
Pyrexia	5 (9.6)	10 (20.4)	9 (18.0)
Hypokalemia	6 (11.5)	13 (26.5)	4 (8.0)
Decreased appetite	5 (9.6)	10 (20.4)	6 (12.0)
Grade 3/4 AEs occurring in ≥10% of patients in any treatment arm, n (%)			
Neutropenia	13 (25.0)	29 (59.2)	14 (28.0)
Thrombocytopenia	4 (7.7)	22 (44.9)	3 (6.0)
Anemia	3 (5.8)	7 (14.3)	1 (2.0)
Leukopenia	2 (3.8)	5 (10.2)	2 (4.0)
Febrile neutropenia	0	6 (12.2)	3 (6.0)
Vomiting	0	5 (10.2)	0
Hypokalemia	1 (1.9)	6 (12.2)	1 (2.0)
SAEs occurring in ≥3% of patients in any treatment arm, n (%)			
Febrile neutropenia	0	6 (12.2)	3 (6.0)
Pneumonia	2 (3.8)	3 (6.1)	0
<i>Pneumocystis jirovecii</i> pneumonia	0	3 (6.1)	0
Myelodysplastic syndrome	0	3 (6.1)	0
Lung infection	0	2 (4.1)	1 (2.0)
Cellulitis	0	0	2 (4.0)
Increased blood lactate dehydrogenase	2 (3.8)	0	0

SAE, serious AE.

investigation of predictive markers of VEN efficacy in FL is warranted.

Acknowledgments

Special thanks to the patients and their families, INVs, study coordinators and support staff, and Genentech CONTRALTO study team members.

Venetoclax is being developed in collaboration between Genentech and AbbVie.

Genentech and AbbVie provided financial support for the study and participated in the design, study conduct, analysis, and interpretation of data, as well as the writing, review, and approval of the presentation. Medical writing support was provided by Catherine Amey and Kate

Authorship

Contribution: All authors were involved with the acquisition of the data; contributed to the analysis and interpretation of the data; provided administrative, technical, or material support or were involved with study supervision; and participated in manuscript development and final approval.

Conflict-of-interest disclosure: P.L.Z. received honoraria, consulting/advisory, and speakers' bureau compensation from Servier, MSD, Gilead, BMS, Janssen, Verastem, and F. Hoffmann-La Roche. I.W.F. received consulting/advisory compensation from AbbVie, Seattle Genetics, TG Therapeutics, and Verastem and research funding from Acerta Pharma, Agios, Calithera Biosciences, Celgene, Constellation Pharmaceuticals, Genentech, Gilead Sciences, Incyte, Infinity Pharmaceuticals, Janssen, Karyopharm Therapeutics, Kile Pharma, Novartis, Pharmacocyclics, Portola Pharmaceuticals, F. Hoffmann-La Roche, TG Therapeutics, Trillium Therapeutics, AbbVie, ArQule, BeiGene, Curis, FORMA Therapeutics, Forty Seven, Merck, Pfizer, Takeda, Teva, Verastem, Gilead sciences, Astra Zeneca, Juno Therapeutics, Unum Therapeutics, and MorphoSys, AG. S.L.S.Y. received travel assistance from Seattle Genetics. M.S.T. received honoraria from Amgen and Regeneron Pharmaceuticals; is on the board of directors or advisory committee of Amgen and F. Hoffmann-La Roche; received research funding from Amgen, Boehringer Ingelheim, Regeneron Pharmaceuticals, and F. Hoffmann-La Roche; and received travel assistance from Amgen. C.R. received honoraria from Celgene, Takeda, Janssen, and F. Hoffmann-La Roche; research funding from Celgene; and travel, accommodation, and expenses assistance from Takeda, F. Hoffmann-La Roche, and Celgene. I.F. received consultancy payments from Novartis, Janssen, Seattle Genetics, Gilead, AbbVie, F. Hoffmann-La Roche, and Celgene and travel, accommodation, and expenses assistance from Gilead, Novartis, and Janssen. C.A. received honoraria and consulting/advisory compensation from Novartis) and travel, accommodation, and expenses assistance from F. Hoffmann-La Roche. G.G. had a consultancy and advisory role at Autolus, Biovista, Takeda, and Italfarmaco; received speakers' bureau compensation from Amgen; received research funding from Gilead; and received travel, accommodation, and expenses assistance from F. Hoffmann-La Roche, Beckton Dickinson, Takeda, and Kite. M.C. received honoraria and consultancy and advisory compensation from F. Hoffmann-La Roche, Servier, Kite, and Gilead. A.P. received employment, travel, accommodation, and expenses compensation from and has ownership

interests in AbbVie. D.S. and E.A.P. are employed by Genentech. D.S. has ownership interests in Genentech. E.A.P. has equity ownership in F. Hoffmann-La Roche and received travel, accommodation, and expenses assistance from F. Hoffmann-La Roche. M.M. has ownership interests and is a former employee of F. Hoffmann-La Roche; has ownership interests in and leadership and employment roles at Corvus Pharmaceuticals. A.S., N.S., M.K., and K.H. are employed by F. Hoffmann-La Roche. E.S.-G. is employed by Genentech and has ownership interests in Genentech/F. Hoffmann-La Roche. M.K. has ownership interests in F. Hoffmann-La Roche, Bayer AG, and Johnson & Johnson. K.H. has equity ownership at F. Hoffmann-La Roche. W.H. received honoraria from F. Hoffmann-La Roche, Gilead, Janssen, and Helsinn Pharmaceuticals; received consultancy payments from F. Hoffmann-La Roche, Gilead, Janssen; is on the speakers' bureau of F. Hoffmann-La Roche, Gilead, Janssen, and Vector; and research funding from F. Hoffmann-La Roche, Janssen, and Amgen. The remaining authors have no competing financial interests.

Correspondence: Wolfgang Hiddemann, Department of Medicine III, University of Munich, Marchioninstr. 15, 81377 Munich, Germany; e-mail: wolfgang.hiddemann@med.uni-muenchen.de.

Footnotes

Submitted 4 March 2020; accepted 13 July 2020; prepublished online on *Blood* First Edition 12 August 2020. DOI 10.1182/blood.2020005588.

Qualified researchers may request access to individual patient level data through the clinical study data request platform (<https://vivli.org/>). Further details on Roche's criteria for eligible studies are available here (<https://vivli.org/members/ourmembers/>). For further details on Roche's Global Policy on the Sharing of Clinical Information and how to request access to related clinical study documents, see https://www.roche.com/research_and_development/who_we_are_how_we_work/clinical_trials/our_commitment_to_data_sharing.htm.

The online version of this article contains a data supplement.

There is a *Blood* Commentary on this article in this issue.

The publication costs of this article were defrayed in part by page charge payment. Therefore, and solely to indicate this fact, this article is hereby marked "advertisement" in accordance with 18 USC section 1734.

REFERENCES

- Czuczman MS, Grillo-López AJ, White CA, et al. Treatment of patients with low-grade B-cell lymphoma with the combination of chimeric anti-CD20 monoclonal antibody and CHOP chemotherapy. *J Clin Oncol*. 1999; 17(1):268-276.
- Herold M, Haas A, Srock S, et al; East German Study Group Hematology and Oncology Study. Rituximab added to first-line mitoxantrone, chlorambucil, and prednisolone chemotherapy followed by interferon maintenance prolongs survival in patients with advanced follicular lymphoma: an East German Study Group Hematology and Oncology Study. *J Clin Oncol*. 2007;25(15):1986-1992.
- Hiddemann W, Kneba M, Dreyling M, et al. Frontline therapy with rituximab added to the combination of cyclophosphamide, doxorubicin, vincristine, and prednisone (CHOP) significantly improves the outcome for patients with advanced-stage follicular lymphoma compared with therapy with CHOP alone: results of a prospective randomized study of the German Low-Grade Lymphoma Study Group. *Blood*. 2005;106(12):3725-3732.
- Marcus R, Imrie K, Solal-Celigny P, et al. Phase III study of R-CVP compared with cyclophosphamide, vincristine, and prednisone alone in patients with previously untreated advanced follicular lymphoma. *J Clin Oncol*. 2008; 26(28):4579-4586.
- Salles G, Mounier N, de Guibert S, et al. Rituximab combined with chemotherapy and interferon in follicular lymphoma patients: results of the GELA-GOELAMS FL2000 study. *Blood*. 2008;112(13):4824-4831.
- Montoto S, López-Guillermo A, Ferrer A, et al. Survival after progression in patients with follicular lymphoma: analysis of prognostic factors. *Ann Oncol*. 2002;13(4):523-530.
- Gallagher CJ, Gregory WM, Jones AE, et al. Follicular lymphoma: prognostic factors for response and survival. *J Clin Oncol*. 1986; 4(10):1470-1480.
- Itchaki G, Brown JR. The potential of venetoclax (ABT-199) in chronic lymphocytic leukemia. *Ther Adv Hematol*. 2016;7(5):270-287.
- Ruefli-Brasse A, Reed JC. Therapeutics targeting Bcl-2 in hematological malignancies. *Biochem J*. 2017;474(21):3643-3657.
- Souers AJ, Levenson JD, Boghaert ER, et al. ABT-199, a potent and selective BCL-2 inhibitor, achieves antitumor activity while sparing platelets. *Nat Med*. 2013;19(2):202-208.
- European Medicines Agency. Venclxyto (venetoclax). Summary of Product Characteristics, Vol. 2019. https://ec.europa.eu/health/documents/community-register/2016/20161205136381/anx_136381_en.pdf. Accessed 16 January 2020.
- US Food and Drug Administration. Full prescribing information for Venclxeta. https://www.accessdata.fda.gov/drugsatfda_docs/label/2019/208573s013lbl.pdf. Accessed 17 December 2019.
- de Vos S, Swinnen LJ, Wang D, et al. Venetoclax, bendamustine, and rituximab in patients with relapsed or refractory NHL: a phase Ib dose-finding study. *Ann Oncol*. 2018;29(9):1932-1938.
- Davids MS, Roberts AW, Seymour JF, et al. Phase I first-in-human study of venetoclax in patients with relapsed or refractory Non-Hodgkin lymphoma. *J Clin Oncol*. 2017;35(8):826-833.

15. Brice P, Bastion Y, Lepage E, et al. Comparison in low-tumor-burden follicular lymphomas between an initial no-treatment policy, prednimustine, or interferon alfa: a randomized study from the Groupe d'Etude des Lymphomes Folliculaires. *Groupe d'Etude des Lymphomes de l'Adulte. J Clin Oncol.* 1997;15(3):1110-1117.
16. Cairo MS, Bishop M. Tumour lysis syndrome: new therapeutic strategies and classification. *Br J Haematol.* 2004;127(1):3-11.
17. World Medical Association. World Medical Association Declaration of Helsinki: ethical principles for medical research involving human subjects. *JAMA.* 2013;310(20):2191-2194.
18. Cheson BD, Fisher RI, Barrington SF, et al; United Kingdom National Cancer Research Institute. Recommendations for initial evaluation, staging, and response assessment of Hodgkin and non-Hodgkin lymphoma: the Lugano classification. *J Clin Oncol.* 2014;32(27):3059-3068.
19. Clopper CJ, Pearson ES. The use of confidence or fiducial limits illustrated in the case of the binomial. *Biometrika.* 1934;26(4):404-413.
20. BA AAC. Approximate is better than "exact" for interval estimation of binomial proportions. *Am Stat.* 1998;52(2):119-126.
21. Kaplan EL, Meier P. Nonparametric estimation from incomplete observations. *J Am Stat Assoc.* 1958;53(282):457-481.
22. Rummel MJ, Al-Batran SE, Kim SZ, et al. Bendamustine plus rituximab is effective and has a favorable toxicity profile in the treatment of mantle cell and low-grade non-Hodgkin's lymphoma. *J Clin Oncol.* 2005;23(15):3383-3389.
23. Robinson KS, Williams ME, van der Jagt RH, et al. Phase II multicenter study of bendamustine plus rituximab in patients with relapsed indolent B-cell and mantle cell non-Hodgkin's lymphoma. *J Clin Oncol.* 2008;26(27):4473-4479.
24. Luminari S, Goldaniga M, Cesaretti M, et al. A phase II study of bendamustine in combination with rituximab as initial treatment for patients with indolent non-follicular non-Hodgkin lymphoma. *Leuk Lymphoma.* 2016;57(4):880-887.
25. Rummel M, Kaiser U, Balsec C, et al; Study Group Indolent Lymphomas. Bendamustine plus rituximab vs fludarabine plus rituximab for patients with relapsed indolent and mantle-cell lymphomas: a multicentre, randomised, open-label, non-inferiority phase 3 trial. *Lancet Oncol.* 2016;17(1):57-66.
26. Seymour JF, Davids MS, Roberts AW, et al. Safety profile of venetoclax monotherapy in patients with chronic lymphocytic leukemia. *Blood.* 2016;128(22):4395.
27. Seymour JF, Ma S, Brander DM, et al. Venetoclax plus rituximab in relapsed or refractory chronic lymphocytic leukaemia: a phase 1b study. *Lancet Oncol.* 2017;18(2):230-240.
28. Roberts AW, Davids MS, Pagel JM, et al. Targeting BCL2 with venetoclax in relapsed chronic lymphocytic leukemia. *N Engl J Med.* 2016;374(4):311-322.
29. Konopleva M, Pollyea DA, Potluri J, et al. Efficacy and biological correlates of response in a phase II study of venetoclax monotherapy in patients with acute myelogenous leukemia. *Cancer Discov.* 2016;6(10):1106-1117.
30. Zelenetz AD, Salles G, Mason KD, et al. Venetoclax plus R- or G-CHOP in non-Hodgkin lymphoma: results from the CAVALLI phase 1b trial. *Blood.* 2019;133(18):1964-1976.
31. Franck Morschhauser PF, Ian W. Venetoclax plus rituximab, cyclophosphamide, doxorubicin, vincristine and prednisolone (R-CHOP) improves outcomes in BCL2-positive first-line diffuse large B-cell lymphoma (DLBCL): first safety, efficacy and biomarker analyses from the phase II CAVALLI study [abstract]. *Blood.* 2018;132(suppl 1). Abstract 782.
32. Vitolo U, Trněný M, Belada D, et al. Obinutuzumab or rituximab plus cyclophosphamide, doxorubicin, vincristine, and prednisone in previously untreated diffuse large B-cell lymphoma. *J Clin Oncol.* 2017;35(31):3529-3537.